

CASE REPORTS

has been reported in 40 percent of male contacts of females with gonorrhea,¹⁸ aggressive tracing of contacts is indicated.

Summary

The Fitz-Hugh—Curtis syndrome, gonococcal peritonitis of the upper abdomen, usually presents with right upper quadrant pain and has rarely been reported in adolescents. The case of a 16-year-old girl in whom there was left upper quadrant pain and in whom initial findings on pelvic examination were normal is described.

REFERENCES

1. Whitwam, PR, Rippberger MS, Shuck JM: Gonococcal perihepatitis presenting as blunt abdominal trauma. *Rocky Mt Med J* 72:383-385, Sep 1975
2. McLain LG, Decker M, Nye D, et al: Gonococcal perihepatitis in a female adolescent: Fitz-Hugh—Curtis syndrome. *JAMA* 239:339-340, Jan 23, 1978
3. Donowitz LG, Lohr JA: Gonococcal disease presenting as right upper quadrant pain (Fitz-Hugh—Curtis syndrome). *Clin Ped* 17:295-296, Mar 1978
4. Stajano C: La reacción frénica en ginecología. *Semana méd* 27:243-248, Aug 19, 1920

5. Claudian I, Florian I: Le syndrome abdominal supérieur et pleurodiaphragmatique aigu au cours annexites gonococciques. *Ann de méd* 43:62-72, Jan 1938
6. Curtis AH: A cause of adhesions in the right upper quadrant. *JAMA* 94:1221-1222, Apr 19, 1930
7. Fitz-Hugh T: Acute gonococcal peritonitis of the right upper quadrant in women. *JAMA* 102:2094-2096, Jun 23, 1934
8. Stanley MM: Gonococcal peritonitis of the upper part of the abdomen in young women. *Arch Intern Med* 78:1-13, Jul 1946
9. Vickers FN, Maloney PJ: Gonococcal perihepatitis: Report of three cases with comments on diagnosis and treatment. *Arch Intern Med* 114:120-123, Jul 1964
10. Trimble C: Gonococcal perihepatitis simulating acute cholecystitis. *Surg Gynecol Obstet* 130:54-56, Jan 1970
11. Kimball MW, Knee S: Gonococcal perihepatitis in a male: The Fitz-Hugh—Curtis syndrome. *N Engl J Med* 282:1082-1084, May 7, 1970
12. Lassus A, Kousa M: Gonococcal perihepatitis and gonococcal caecitis—Presentation of a case with cutaneous manifestations. *Br J Vener Dis* 49:48-49, Feb 1973
13. Reichert JA, Valle RF: Fitz-Hugh—Curtis syndrome: A laparoscopic approach. *JAMA* 236:266-268, Jul 19, 1976
14. Gästrin B, Kallings LO: Improved methods for gonococcal sampling and examination on a large scale. *Acta Pathol Microbiol Scand* 74:362-370, 1968
15. Wald E: Gonorrhea—Diagnosis by gram stain in the female adolescent. *Am J Dis Child* 131:1094-1096, Oct 1977
16. Holmes KK: Gonococcal infection: Clinical, epidemiologic and laboratory perspectives. *Adv Intern Med* 19:259-285, 1974
17. Gonorrhea—CDC recommended treatment schedules. *Morbidity Mortality Weekly Rep* 23:341-348, Oct 5, 1974
18. Handsfield HH, Lipman TO, Harnisch JP, et al: Asymptomatic gonorrhea in men: Diagnosis, natural course, prevalence and significance. *N Engl J Med* 290:117-123, Jan 17, 1974

Refer to: Liggett C, Kartchner M: Peripheral arterial tumor embolism by malignant tumor. *West J Med* 130:72-75, Jan 1979

Peripheral Arterial Tumor Embolism by Malignant Tumor

CHARLES LIGGETT, MD
MARK KARTCHNER, MD
Tucson, Arizona

EMBOLIZATION OF A peripheral artery most commonly originates from the heart. Mural thrombus secondary to atrial fibrillation and myocardial infarction is the usual cause.^{1,2} Rarer etiologies include atrial myxoma,³ paradoxical embolism,^{4,5} atherosclerosis,⁶⁻⁹ subacute bacterial endocarditis,¹⁰ fungal endocarditis,¹¹ release of prosthetic valve material,¹² idiopathic hypertrophic subaortic stenosis,¹³ hydatid cyst,¹⁴ echinococcal cyst,¹⁵ ventricular aneurysms,¹⁶ bullets,^{17,18} mitral stenosis,^{19,20} false aneurysms²¹ and arterial malignancy.²² This paper presents the 32nd reported case of tumor embolism and reviews its salient features.

From the Department of Surgery, Tucson Hospitals Medical Education Program, Tucson, Arizona.

Submitted, revised, April 17, 1978.

Reprint requests to: Mark M. Kartchner, MD, 5402 East Grant Road, Suite B5, Tucson, AZ 85712.

Report of a Case

A 53-year-old woman was admitted to Tucson Medical Center on January 21, 1977, with a 2½-hour history of severe pain, numbness and coldness of both legs. She had a history of organic heart disease and hypertension, but she was asymptomatic in those respects. She had been followed for the past year for a nonexpanding nodule in her right lung (Figure 1). On physical examination there were no pulses in the right groin or in distal arteries of either leg. A weak left femoral pulse was palpable.

An immediate retrograde bilateral aortoiliac and femoral embolectomy was carried out through common femoral arteriotomies and good flow was established in the extremities. The pathology report showed poorly differentiated carcinoma in the clot.

Tomography showed that the pulmonary mass was larger than was shown on routine x-ray of the chest. Examination of subsequent bronchoscopic washings showed possibility of malignancy. On February 9, 1977, an exploratory thoracotomy and right lower lobectomy were done. Postoperatively she was unresponsive because of cerebral vascular insult; she died February 12, 1977. On postmortem examination, an undifferentiated sarcoma of unknown cause was found with multiple tumor emboli to the right kidney, brain and liver.

CASE REPORTS

The pulmonary lesion was a metastatic sarcoma (Figures 2, 3).

Discussion

Tumor is a rare cause of arterial embolization, and this is the ninth reported case of successful removal of a malignant tumor embolus (Table 1). Pathologic examination of the embolic specimen led to the correct diagnosis, illustrating the need for pathologic examination of all embolic material. All but one of the reported cases of tumor embolism were associated with lung malignancy, and the remaining one was due to aortic invasion of a pelvic osteogenic sarcoma with

distal embolization.²² The occurrence of tumor embolization during surgical operation for pulmonary cancer has been reported by others (Table 2), and several authors have suggested obtaining early venous control in resecting large lung tumors with known venous invasion.^{6,12,29,30,32,36}

Summary

A rare case of peripheral arterial embolization by malignant tumor is reported and the literature




Figure 1.—Admission x-ray of chest showing lesion below right inferior pulmonary artery.



Figure 2.—Low-power view of tumor embolus in pulmonary vein.




Figure 3.—High-power view showing tumor embolus in pulmonary vein.

TABLE 1.—Systemic Tumor Embolism, Successful Removal

<i>Authors and Year of Publication</i>	<i>Origin of Tumor Embolus</i>	<i>Obstructed Arteries</i>
Mital and associates, 1971 ²³	Thyroid malignancy with pulmonary metastasis	Aortic bifurcation
DeBoer and Prillelits, 1969 ²⁴	Undifferentiated lung carcinoma*	Femoral artery
Groth, 1940 ²⁵	Tibial sarcoma with lung metastasis	Femoral artery
Blum, 1950 ²⁶	Oat cell carcinoma of lung	Femoral artery
Firor and Pearson, 1967 ²⁷	Squamous cell carcinoma of lung*	Aortic bifurcation
Soriano and associates, 1964 ²⁸	Osteogenic sarcoma of fibula with lung metastasis	Femoral artery
MacMahon and associates, 1974 ²⁹	Giant cell carcinoma of lung*	Distal aorta
	Undifferentiated carcinoma of lung*	Brachial artery
Personal observation	Metastatic sarcoma to lung*	Aortic bifurcation

*See Table 2.

CASE REPORTS

is reviewed. Tumor emboli are usually pulmonary in origin, and they are commonly associated with surgical operation for pulmonary cancer. The value of pathologic and bacteriologic examination of all embolic material is emphasized.

REFERENCES

1. Darling RC, Austen WG, Linton RR: Arterial embolism. *Surg Gynecol Obstet* 124:106-114, Jan 1967
2. Thompson JE, Weston AS, Sigler L, et al: Arterial embolism after acute myocardial infarction. *Ann Surg* 171:979-985, Jun 1970
3. Thompson JR, Simmons CR: Arterial embolus: Manifestation of unsuspected myxoma. *JAMA* 288:864-865, May 13, 1974
4. Davies DJ, Muller HK, Cooper E: Paradoxical embolism. *Med J Aust* 2:1120-1121, Dec 1973
5. Paradoxical embolism (Editorial). *Br Med J* 1:630-631, Mar 1963
6. Aylwin JA: Avoidable vascular spread in resection for bronchial carcinoma. *Thorax* 6:250-267, Sep 1951
7. Kwaan JHM, Molen RU, Stemmer EA, et al: Peripheral embolism resulting from unsuspected atheromatous aortic plaques. *Surgery* 78:583-587, Nov 1975
8. Lord JW, Rossi G, Daliana M, et al: Unsuspected abdominal aortic aneurysms as the cause of peripheral arterial occlusive disease. *Ann Surg* 177:767-771, Jun 1973
9. Mehigan JT, Stony RJ: Lower extremity atheromatous embolization. *Am J Surg* 132:163-167, Aug 1976
10. Gustke R, Morgan D: Five years experience with bacterial endocarditis at the West Virginia University Medical Center. *Dis Chest* 50:265-269, Sep 1966
11. Montague NT, Sugg WL: Candida endocarditis with femoral emboli. *J Thorac Cardiovasc Surg* 67:322-327, Feb 1974
12. Pluth JR, Danielson GK: Poppet embolism in cloth-covered silastic poppet valves. *Mayo Clinic Proc* 49:811-814, Nov 1974
13. Kawarda Y, Najmi M, Matsumoto T: A case report of arterial emboli in a male secondary to idiopathic hypertrophic subaortic stenosis. *Vasc Surg* 7:276-283, Nov-Dec 1973
14. Aletras HA: Embolism of the popliteal artery by a ruptured cyst of the heart. *Scand J Thorac Cardiovasc Surg* 1:188-190, 1967
15. Christeas N, Balas P, Chadjiyiannakin E: Embolism of the femoral artery by an echinococcus cyst successfully treated by embolectomy. *Am J Surg* 115:673-674, May 1968
16. Chesler MB, Tucker RBK, Barlow MB: Subvalvular and apical left ventricular aneurysms in the Bantu as a source of systemic emboli. *Circulation* 35:1156-1161, Jun 1967
17. DiMaio VJ, DiMaio DJ: Bullet embolism. *J Forensic Sci* 17:394-398, Jul 1972
18. Rodriguez MA, Rodger MR: Right axillary artery embolus following gunshot wound of the back. *J Trauma* 15:170-173, Feb 1975
19. Casella L, Abelman WH, Ellis LB: Patients with mitral stenosis and systemic emboli. *Arch Intern Med* 114:773-781, Dec 1964
20. Lillicrap D, Piersowicz A: Mitral stenosis and systemic emboli. *Br Med J* 2:1169-1171, Nov 1964
21. Waibel PP, Lundin H: Unusual source for peripheral arterial embolization. *Arch Surg* 92:105-106, Jan 1966
22. Till AS, Fairburn EA: Massive neoplastic embolism. *Br J Surg* 35:86-89, Jul 1947
23. Mital RN, Prakash A, Goel RG: Undifferentiated malignant tumor of the thyroid gland with tumor embolism of the aorta. *J Cardiovasc Surg* 12:75-77, Jan-Feb 1971
24. DeBoer HH, Prillevis HW: Massive tumor embolism. *Arch Chir Neerl* 21:223-234, 1969
25. Groth KE: Tumor embolism of the common femoral artery treated by embolectomy and heparin. *Surgery* 8:617-631, Oct 1940
26. Blum L: Successful removal of a tumor embolus from the femoral artery. *JAMA* 142:981-987, Apr 1, 1950

TABLE 2.—Postthoracotomy Tumor Embolism

<i>Authors and Year of Publication</i>	<i>Primary Tumor</i>	<i>Obstructed Arteries</i>
DeBoer and Prillevis, 1969 ²⁴	Undifferentiated lung carcinoma	Femoral artery
Balas and associates, 1971 ³⁰	Undifferentiated bronchial carcinoma	Aortic bifurcation
Probert, 1956 ³¹	Adrenocortical carcinoma with lung metastasis	Common carotid and innominate arteries
Eason, 1950 ³²	Squamous cell carcinoma of lung	Cerebral arteries
Aylwin, 1951 ⁶	Bronchial carcinoma	Internal and axillary arteries
Taber, 1961 ³³	Fibrosarcoma with lung metastasis	Aorta
	Pulmonary adenocarcinoma	Aortic bifurcation and multiple organ arteries
Christiansen and Morgan, 1965 ³⁴	Undifferentiated lung carcinoma	Aortic bifurcation
Senderoff and Kirschner, 1962 ³⁵	Thyroid carcinoma with lung metastasis	Femoral artery
Firor and Pearson, 1967 ²⁷	Squamous cell carcinoma	Aortic bifurcation
MacMahon and associates, 1974 ²⁹	Giant cell carcinoma of lung	Aortic bifurcation
	Undifferentiated carcinoma of lung	Brachial artery
Personal observation	Metastatic sarcoma to lung	Multiple organ arteries

TABLE 3.—Systemic Tumor Embolism, Unsuccessful Removal or Diagnosed at Autopsy

<i>Authors and Year of Publication</i>	<i>Origin of Tumor Embolus</i>	<i>Obstructed Arteries</i>
Rudusky, 1965, cited by		
DeBoer and Prillevis, 1969 ²⁴	Maxillary chondrosarcoma with lung metastasis	Aorta and iliac artery
Loosemore and Whittick, 1951 ³⁷	Synovioma of knee with lung metastasis	Axillary artery
Buckmaster, 1961 ³⁸	Oat cell carcinoma of lung	Aortic bifurcation
Green and associates, 1974 ³⁹	Colonic adenocarcinoma with lung metastasis	Multiple organ arteries
Till and Fairburn, 1947 ²²	Oat cell carcinoma of lung	Femoral artery
Cera and associates, 1957 ⁴⁰	Anaplastic bronchogenic carcinoma with atrial invasion	Coronary artery
Busse, 1903 ⁴¹	Chorioepithelioma metastatic to lung	Cerebral embolism
Storjohann, 1932 ³⁶	Chorioepithelioma metastatic to lung	Cerebral embolism
Kaufmann, 1929, cited by		
Loosemore and Whittick ³⁷	Chorionic carcinoma with lung metastasis	Splenic and renal arteries
Miller and Jackson, 1954 ⁴²	Pulmonary myxosarcoma	Splenic and cerebral arteries
Van Way and Lawler, 1969 ⁴³	Osteogenic sarcoma involving the pelvis and distal aorta	Femoral arteries
Lisa and associates, 1941 ⁴⁴	Squamous cell carcinoma of lung	Coronary artery
Hiraoka and associates, 1976 ⁴⁵	Breast carcinoma metastatic to lung	Coronary artery
Haiby and Baiton, 1965 ⁴⁶	Bronchial adenoma	Multiple organ arteries
Mori, 1964, cited by		
Hiraoka and associates ⁴⁵	Lung cancer	Coronary artery

CASE REPORTS

27. Firor WB, Pearson FG: Massive systemic tumor embolism during pneumonectomy: Successful removal by means of balloon catheter. *Can J Surg* 10:200-202, Apr 1967
28. Soriano FM, Berry RW, Shen JP: Experiences with arterial tumor embolism. *St. Vincent's Hosp Med Bul* 6:33-35, Jun 1964
29. MacMahon H, Forrest JV, Weisz D, et al: Embolism occurring during pneumonectomy. *Ann Thorac Surg* 17:395-397, Apr 1974
30. Balas P, Katsaras E, Zoiopoulos M: Peripheral arterial embolism by malignant tumor. *Vasc Surg* 5:27-29, Jan-Feb 1971
31. Probert WR: Sudden operative death due to massive tumour embolism. *Br Med J* 1:435-436, Feb 1956
32. Eason EH: A case of cerebral infarction due to neoplastic embolism. *J Pathol* 62:454-457, Jul 1950
33. Taber RE: Massive systemic tumor embolization during pneumonectomy. *Ann Surg* 154:263-268, Aug 1961
34. Christiansen TW, Morgan S: Tumor embolism in a peripheral artery. *Ann Thorac Surg* 1:311-313, May 1965
35. Senderoff E, Kirschner PA: Massive tumor embolism during pulmonary surgery. *J Thorac Cardiovasc Surg* 44:528-535, Oct 1962
36. Storjohann KR: Ein Fall von Chorianepitheliom im Hoden mit Gynakomastie. *Frankfort Ztschr Path* 43:80-95, 1932
37. Loosemore TGE, Whittick JW: Gross tumour embolism of the axillary artery by malignant synoviomia. *J Pathol* 63:133-138, Jan 1951
38. Buckmaster NS: Simultaneous bilateral tumor embolism of the common iliae arteries. *Med J Aust* 2:516-518, Sep 1961
39. Green WH, Benjamin RS, Glusmar S, et al: Arterial embolism of tumor causing fatal organ infarction. *Arch Intern Med* 134:545-548, Sep 1974
40. Cera LJ, Karlinsky W, Rodin AE: Tumor embolism of the left coronary artery. *Am Heart J* 53:472-478, Mar 1957
41. Busse O: Ueber Chorioepitheliome, die ausserhalb der Placentarsfelle entstanden sind. *Arch Pathol Anat Physiol Clin Med* 174:207-231, 1903
42. Miller AA, Jackson FB: Gross arterial embolism by a myxosarcoma of pulmonary origin. *J Pathol* 68:221-226, Jul 1954
43. Van Way CW III, Lawler MR: Osteogenic sarcomatous emboli to the femoral arteries. *Am J Surg* 117:745-747, May 1969
44. Lisa JR, Hirschhorn L, Hart CA: Tumors of the heart. *Arch Intern Med* 67:91-113, Jan 1941
45. Hiraoka K, Ohkawa S, Uedu W, et al: A case of tumor embolism of the coronary artery resulting in myocardial infarction and cardiac rupture. *Jap Heart J* 17:269-274, Mar 1976
46. Haiby G, Baiton D: Tumor embolism to coronary artery after pneumonectomy. *JAMA* 192:256-257, Apr 19, 1965

Refer to: Edwards DA, Davis PR Jr, Johnson MC, et al: Delayed fat embolism after total hip arthroplasty. *West J Med* 130:75-77, Jan 1979

Delayed Fat Embolism After Total Hip Arthroplasty

DAVID A. EDWARDS, MD
PHILLIP R. DAVIS, JR., MD
MICHAEL C. JOHNSON, MD
WILLIAM A. TEIPNER, MD
Reno, Nevada

THE SYNDROME of fat embolism is a serious clinical entity, the pathogenesis of which remains obscure. Although several reports have appeared describing the onset of the syndrome after joint replacement using methylmethacrylate, delayed onset of the syndrome has not been reported.¹ The present case report deals with a patient in whom this syndrome developed with rather late onset and recurrent features. The possible rela-

From the Departments of Medicine (Drs. Edwards, Davis and Johnson), and Orthopedic Surgery (Dr. Teipner), St. Mary's Hospital, Reno, Nevada.

Submitted March 27, 1978.

Reprint requests to: David A. Edwards, MD, Internal Medicine Associates, 175 West 6th Street, Suite 23, Reno, NV 89503.

tionship to the use of methylmethacrylate will be discussed.

Report of a Case

A 64-year-old previously healthy man was initially seen because of increasing pain in the right hip. Vital signs were normal with the exception of an irregular pulse. There were bilateral rhonchi and wheezes; four to eight extrasystoles per minute were noted. An electrocardiogram showed unifocal premature ventricular contractions (PVC) and complete left bundle branch block. Results of complete blood count, analysis of urine and SMA-12 were normal. An x-ray film of the hips showed severe degenerative joint disease involving the right hip. The preoperative roentgenogram of the chest is shown in Figure 1.

A total hip arthroplasty was done on February 18, 1977, using an Aufranc-Turner prosthesis. The prosthesis was held in place with methylmethacrylate cement. The duration of the surgical operation was 110 minutes. The patient's vital signs were stable throughout the procedure. His urine output remained excellent during operation, exceeding 100 ml per hour, and the total blood loss was estimated at 1,200 ml. He was given 3,000 ml of lactated Ringer's solution and 1,000 ml of whole blood during the procedure. Continuous electrocardiographic monitoring showed rare PVC. Recovery in the immediate postoperative period was uneventful and the patient was transferred back to the ward in excellent condition. Postoperative treatment consisted of administration of cephalothin sodium, 1 gram given every four hours; heparin sodium, 2,000 units given intravenously every six hours, and morphine sulphate, given as needed for pain.

On the morning following operation the patient was resting comfortably. His temperature was 38°C (100.4°F) and other vital signs were normal. Total output was 3,130 ml (2,920 ml as urine and 210 ml from wound drainage). The hemoglobin and hematocrit values were stable at 13.3 grams per dl and 38.7 percent, respectively. Findings on an electrocardiogram (ECG) remained unchanged.

Thirty-six hours postoperatively the patient's temperature rose abruptly to 39.2°C (102.5°F), his respiratory rate was 36 breaths per minute and pulse was 120 beats per minute. He was confused, restless, disoriented, and incoherent. Diffuse rales were heard in both lungs and a petechial rash in the axillary and hypogastric regions de-